


Advanced Electrodeless Plasma Propulsion Concepts



Edgar Choueiri


Electric Propulsion & Plasma Dynamics Lab
Princeton University

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 25 JUN 2004		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Advanced Electrodeless Plasma Propulsion Concepts				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Electric Propulsion & Plasma Dynamics Lab Princeton University				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001793, International Symposium on Energy Conversion Fundamentals Held in Istanbul, Turkey on 21-25 June 2005., The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 35	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Outline

- I. FARAD: Faraday Accelerator with RF-Assisted Discharge
 - Basic Concept
 - Advantages
 - Basic Questions and Approach
 - Results of Proof-of-concept Experiment
- II. Ion Acceleration by Beating Electrostatic Waves
 - Basic Concept
 - Advantages
 - Basic Challenges
 - Status of Research

FARAD: Faraday Accelerator with RF-Assisted Discharge




FARAD: Advantages

General:

- Completely electrodeless
- EM acceleration: High thrust density & I_{sp}
- Compatibility with a wide variety of propellants
- High mass utilization efficiency
- Accelerating forces are *always* perpendicular to \mathbf{B} : No detachment issues
- Pulsed: can be used on power-limited s/c

With respect to PIT:

- Much more compact device
- No obstacles in plasma stream
- Additional axial Lorentz force: $J_{\theta} B_r$
- Plasma confinement through $J_{\theta} B_z$ leads to low wall losses



FARAD: Basic Questions


- What are the fundamental features and governing relations of the acceleration mechanisms?
- What are the fundamental limitations and major loss mechanisms?
- What are the scaling laws?
- How to choose the various controllable parameters (propellant, injected mass bit, RF power, B-field strength and topology, RF pulse length, inductive coil current and pulse length, geometrical dimensions, etc.) in order to optimize the performance?

FARAD: Approach


- ✓ • Develop **Proof-of-concept Experiment** and demonstrate current sheet formation and acceleration.
- Extensive Characterization of Plasma, Current Sheet and accel. mechanisms
- Analytical and Numerical Modeling
- Optimization of Operational Parameters
- Design and Manufacturing a Prototype FARAD Thruster
- Extensive Performance Testing and Optimization

FARAD: Proof-of-Concept Experiment




Concept




Experiment





FARAD: Proof-of-Concept Experiment: J_θ coil




FARAD: Proof-of-Concept Experiment: Set-up




Advanced Electrodeless Concepts: Experimental Field



FARAD: Proof-of-Concept Experiment: Demonstration of passive plasma feeding




FARAD: Proof-of-Concept Experiment: Demonstration of Current Sheet Formation and Acceleration



II. Ion Acceleration by Beating Electrostatic Waves




Inspiration : Ionospheric Observations



- Low-energy ionospheric oxygen ions **naturally** accelerated and reach escape velocities.
- Intense Lower Hybrid Waves: ~ 100 mV/m, 2-12 kHz, Electrostatic in nature.
- Classical ion acceleration by (**resonant**) ES waves fails to explain the observations. (Initial ion velocity < Threshold interaction velocity)
- 1998: Benisti et. al. (MIT) propose that a beating of 2 ES waves may be responsible.

J. Geophys. Res. **103** 9431
(1998)

Beating Waves Vs. Single Wave




What are the conditions for Ion acceleration by Beating waves?

Necessary Condition: $\omega_i - \omega_j = n\omega_c$




Resonant Energization

$$\omega = \omega_{ci}$$





Beating Wave Energization

$$\omega_2 - \omega_1 = \omega_{ci}$$



Benisti's Criterion is not Sufficient!

Choueiri & Spektor, AIAA-2000-3759, IEPC-01-209



What are the necessary
and sufficient conditions
for ion acceleration by
beating waves?

Are there Necessary and Sufficient Criteria for Acceleration?




Yes

Spektor & Choueiri, *Physical Review E*, March 2004




Model – Basic Concept

Single particle, constant magnetic field, multiple waves



$$\ddot{x} + \omega_{ci}^2 x = \frac{q}{m} \sum_i E_i \cos(\kappa_i x - \omega_i t)$$

Approach






Poincare diagram construction

1 - D periodic Motion


$$(\rho, \theta, t) = (\rho, \theta, t+T)$$

$T = \text{period}$




Trapped Particles & Domain of Allowed Acceleration

Numerical Simulation



2nd-order Perturbation Theory



Hamilton's Eqns

$$\begin{aligned}\dot{\rho} &= \frac{\partial H}{\partial \theta} = \varepsilon \{ \nu_i J_{\nu_i}(\rho) \sin(\nu_i \theta) \\ &+ \nu_j J_{\nu_j}(\rho) \sin(\nu_j \theta) \} \\ &+ \varepsilon^2 (\nu_i - \nu_j) S_6^{\nu_i, \nu_j}(\rho) \sin[(\nu_j - \nu_i) \theta] = 0, \\ \dot{\theta} &= -\frac{\partial H}{\partial \rho} = \varepsilon \{ J'_{\nu_i}(\rho) \cos(\nu_i \theta) \\ &+ J'_{\nu_j}(\rho) \cos(\nu_j \theta) \} \\ &+ \varepsilon^2 \{ S_1^{\nu_i}(\rho) + S_1^{\nu_j}(\rho) \\ &+ S_6^{\nu_i, \nu_j}(\rho) \cos[(\nu_i - \nu_j) \theta] \} = 0.\end{aligned}$$


$$\begin{aligned}H &= \frac{\varepsilon^2 \pi}{8 \sin \nu \pi} \left(1 + \frac{\rho}{\nu} \cos \theta \right) \\ &\left[- J_{\nu-1}(\rho) J_{-(\nu-1)}(\rho) + J_{\nu}(\rho) J_{-\nu}(\rho) \right. \\ &\quad \left. + J_{\nu+1}(\rho) J_{-(\nu+1)}(\rho) - J_{\nu+2}(\rho) J_{-(\nu+2)}(\rho) \right]\end{aligned}$$

Hamiltonian from 2nd order perturbation theory + Lie Transforms

Bessel functions properties

$$\begin{aligned}F(\rho, \nu) &\simeq \frac{\rho}{\nu} - 1 + \frac{1}{2} \\ &3 \left(f_{\nu-1} + f_{-(\nu-1)} + f_{\nu+2} + f_{-(\nu+2)} \right) = 0\end{aligned}$$

where $f_{\nu} = f_{\nu}(\rho) \equiv J'_{\nu} / J_{\nu}$



Simplified Eqn. for locating critical points

Necessary and Sufficient Criteria

Benisti's beat criterion:

$$\omega_2 - \omega_1 = n\omega_c$$

Additional criterion:


$$H_H < H(\rho_0, \theta_0) < H_E$$

$$H_H = H(\rho \simeq \nu - \sqrt{\epsilon}; \theta = \pi).$$



$$H_H \simeq H(v \simeq \frac{\omega}{k}; \theta = \pi).$$


$$H_E = H(\rho \simeq \frac{\nu - \sqrt{\epsilon}}{2}; \theta = \pi).$$



$$H_E \simeq H(v \simeq \frac{\omega}{2k}; \theta = \pi).$$

Does it work for many particles with collisions?

Monte Carlo simulations



- Ion – ion collisions enhance acceleration!

- Enhancement is due to depopulation of trapped region of phase space due to collisions.

$$v_{ii} \sim 10 \cdot \omega_{ci}$$





Summary of Theoretical Findings

- Previously proposed criterion is necessary but not sufficient.
- Poincare diagram shows that trapped particles lie between critical elliptic and hyperbolic points.
- New criterion found and relies on the value of the Hamiltonian at the motion's critical points.
- Collisions enhance the effect.
- Ions with arbitrarily low initial velocity can be accelerated in contrast with non-beating waves.
 - Seems to occur naturally in the ionosphere
 - Promise for propulsion applications and more efficient ion heating.

TIME FOR EXPERIMENTAL VERIFICATION

Experimental Apparatus



RF Power: 100 – 1200 Watt

Magnetic Field: 0.08 Tesla

Plasma Density: $10^{10} - 10^{13} \text{ cm}^{-3}$

$T_e \sim 3\text{-}5 \text{ eV}$, $T_i \sim 0.1\text{-}0.3 \text{ eV}$

Experimental Apparatus





Base Pressure: $\sim 10^{-6}$ Torr

Operating Pressure: $\sim 10^{-3}$ Torr

Electric Circuit for Helicon Source

P_{RF} up to 1.2 kW



$\omega \sim 13.56$ MHz



Dedicated Diagnostics

- Langmuir Probes (n_e , T_e)
- Hall Probes (B field)
- Probe interferometry (Dispersion Relation)
- Retarding Potential Analyzer (Ion velocity)
- LIF (Ion Velocity)

Launching Waves




Langmuir Probes

- Need to launch an electrostatic wave \perp to the magnetic field
- Measuring the dispersion to determine the nature of the wave

Detecting the Waves

Interferometry



125 W Interferometry Data

Integer/decimal





.2

.4

.6





.8

$v=1.$







$f_{ci}=30$ kHz


$v=2.$



$v=3.$



Internal Antenna – Data Interpretation



- Electrostatic Ion Cyclotron fast wave is launched
- Good agreement with theory
- Indirect evidence of ion energization
- Direct evidence will be obtained next month with LIF

Conclusions

- Two electrodeless plasma acceleration concepts with promised for pulsed and steady-state propulsion are investigated
 - FARAD: Proof of concept experiment verified basic principle
 - Beating wave ion acceleration:
 - Found fundamental acceleration criteria
 - Monte Carlo simulation verified that the effect can exist in a real plasma with collisions
 - Dedicated experiment is yielding first laboratory observations of new acceleration mechanism.